EEF1D Polyclonal Antibody

catalog number: E-AB-19083



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Immunogen Fusion protein of human EEF1D

Host Rabbit Isotype IgG

Purification Antigen affinity purification


Conjugation Unconjugated

buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.


Applications	Recommended Dilution
WB	1:1000-1:5000

IHC 1:50-1:300

Data




Western blot analysis of A549 A172 and HepG2 cell lysates using EEF1D Polyclonal Antibody at dilution of 1:950



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using EEF1D Polyclonal Antibody at dilution of 1:65(×200)

Observed-MV: Refer to figures

Calculated-MV:31 kDa



Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using EEF1D Polyclonal Antibody at dilution of 1:65(×200)

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

EEF1D, also named as EF1D and EF 1 delta, belongs to the EF-1-beta/EF-1-delta family. It is a subunit of the elongation factor-1 complex, which is responsible for the enzymatic delivery of aminoacyl tRNAs to the ribosome. EF-1-beta and EF-1-delta stimulate the exchange of GDP bound to EF-1-alpha to GTP. EEF1D is phosphorylated upon DNA damage, probably by ATM or ATR. This antibody is a mouse monoclonal antibody raised against residues near the C terminus of human EEF1D.

For Research Use Only