

Purified Anti-Mouse CD28 Antibody[37.51], Functional Grade

catalog number: E-AB-F10260

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse
Immunogen	Recombinant Mouse CD28 protein
Host	Syrian Hamster
Isotype	Syrian Hamster IgG
Clone	37.51
Purification	>98%, Protein A/G purified
Buffer	Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method.

Applications

Recommended Dilution

FCM	≤ 0.2 µg per million cells in 100 µL volume
Activ	Reported in the literature
Stim	Reported in the literature


Data



C57/BL6 Mouse splenocytes were stained with 0.2 µg Purified Anti-Mouse CD28 Antibody[37.51], Functional Grade (Right) and 0.2 µg Syrian Hamster IgG2, κ Isotype

Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Syrian Hamster IgG Secondary Antibody, then anti-Mouse CD3 PE-conjugated Monoclonal Antibody.

Data shown was gated on CD3 + cell population.



C57/BL6 Mouse splenocytes were stained with 0.2 µg Purified Anti-Mouse CD28 Antibody[37.51], Functional Grade (Right) and 0.2 µg Syrian Hamster IgG2, κ Isotype

Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Syrian Hamster IgG Secondary Antibody, then anti-Mouse CD3 PE-conjugated Monoclonal Antibody. Data

shown was gated on CD3 + cell population.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.
Shipping	Ice bag

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V2.6

CD28 is a 44 kD glycoprotein, also known as Tp44 or T44. It is a member of the Ig superfamily, expressed on thymocytes, most peripheral T cells, and NK cells. In association with CD80 (B7-1) and CD86 (B7-2), CD28 acts as the second signal for T and NK cell activation and proliferation. The 37.51 antibody has been reported to augment in vitro T cell proliferation and cytokine production, and promote CTL development.

None (Azide-Free, Low Endotoxin) are perfectly suited to be used in culture or in vivo (for nonhuman studies) for functional assays blocking, neutralizing, activation or depletion where the presence of azide may damage cells or exogenous endotoxin may signal or activate cells.

Application References

Ai-Di Gu, et al. Immunity. 2015 Jan 20;42(1):68-79.