Purified Anti-Human/Monkey CD11b Antibody[ICRF44], Functional Grade

catalog number: **E-AB-F11460**


Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Reactivity	Human;Monkey
Immunogen	Recombinant Human/Monkey CD11b protein
Host	Mouse
Isotype	Mouse IgG1, κ
Clone	ICRF44
Purification	> 98%, Protein A/G purified, ≥ 95% monomer by analytical SEC
Endotoxin	< 1 EU/mg determined by LAL assay
Conjugation	Unconjugated
Buffer	Sterile PBS, pH 7.2. Contains no stabilizers or preservatives

Applications	Recommended Dilution
FCM	≤ 0.2 μg per million cells in 100 μL volume
Block	Reported in the literature

Data



Human/Monkey peripheral blood monocytes were stained with 0.2 μg Purified Anti-Human/Monkey CD11b Antibody[ICRF44], Functional Grade (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD14 FITC-conjugated Monoclonal Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze/thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.
Shipping	Ice bag

Background

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com

Human ITGAM (Integrin alpha M subunit) is a 127kDa (predicted) glycoprotein, a member of the Integrin family of proteins. The Integrin family proteins are heterodimeric transmembrane receptors composed of an alpha and a beta subunit. The Integrin alpha M subunit, also known as MAC-1 alpha subunit or CD11b, combines with the Integrin beta 2 subunit (CD18) to form the non-covalent heterodimer Integrin alpha M/ beta 2, also known as MAC-1 and complement receptor type 3 (CR3).

None (Azide-Free, Low Endotoxin) are perfectly suited to be used in culture or in vivo (for nonhuman studies) for functional assays blocking, neutralizing, activation or depletion where the presence of azide may damage cells or exogenous endotoxin may signal or activate cells.

Application References

Sprong T, et al. Blood. 2003;102:3702.