## Recombinant Human ILT4/LILRB2 protein (His Tag)

## Catalog Number: PDMH100418

Note: Centrifuge before opening to ensure complete recovery of vial contents.

SpeciesHumanSourceHEK293 Cells-derived Human ILT4 protein Met1-Val461, with an C-terminal HisCalculated MW50.6 kDaObserved MW75 kDaAccessionQ8N423Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU/mg of the protein as determined by the LAL method	Description			
Calculated MW50.6 kDaObserved MW75 kDaAccessionQ8N423Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU/mg of the protein as determined by the LAL method	Species	Human		
Observed MW75 kDaAccessionQ8N423Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU/mg of the protein as determined by the LAL method	Source	HEK293 Cells-derived Human ILT4 protein Met1-Val461, with an C-terminal His		
AccessionQ8N423Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU/mg of the protein as determined by the LAL method	Calculated MW	50.6 kDa		
Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU/mg of the protein as determined by the LAL method	Observed MW	75 kDa		
Properties   Purity > 95% as determined by reducing SDS-PAGE.   Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method   Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.   Shipping This product is provided as lyophilized powder which is shipped with ice packs.   Formulation Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.   Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of	Accession	Q8N423		
Purity> 95% as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU/mg of the protein as determined by the LAL method	Bio-activity	Not validated for activity		
Endotoxin< 1.0 EU/mg of the protein as determined by the LAL methodStorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.ReconstitutionIt is recommended that sterile water be added to the vial to prepare a stock solution of	Properties			
StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Purity	> 95% as determined by reducing SDS-PAGE.		
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method		
ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.ReconstitutionIt is recommended that sterile water be added to the vial to prepare a stock solution of	Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80		
ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.ReconstitutionIt is recommended that sterile water be added to the vial to prepare a stock solution of		°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.ReconstitutionIt is recommended that sterile water be added to the vial to prepare a stock solution of		reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.		
ReconstitutionMannitol.It is recommended that sterile water be added to the vial to prepare a stock solution of	Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
<b>Reconstitution</b> It is recommended that sterile water be added to the vial to prepare a stock solution of	Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with 5% Trehalose and 5%		
		Mannitol.		
$0.5 \mathrm{mg/mL}$ . Concentration is measured by UV-Vis	Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of		
o.o mg/me. concentration is incustice by 0 v vis.		0.5 mg/mL. Concentration is measured by UV-Vis.		

## Data

М	R
10	-
-	
-	
1	
	M

> 95 % as determined by reducing SDS-PAGE.

## Background

Members of the immunoglobulin-like transcript (ILT) family are activating and inhibitory immunoreceptors whose genes are located same locus that encodes killer cell Ig-like receptors (KIR). Leukocyte Immunoglobulin-Like Receptor Subfamily B Member 2 (LIR-2) is a type I transmembrane protein. LIR-2 is expressed primarily on monocytes and dendritic cells (DC). Human LIR-2 is produced as a 598 amino acino acid precursor including a 21 aa signal sequence, a 440 aa extracellular domain (ECD), a 21 aa transmenbrane segment, and a 116 aa cytoplasmic domain. LIR-2 binds to Classical MHCI proteins. Ligation of LIR-2 incluces Tyr phosphorylation within its cytoplasmic ITIMs, a requirement for association with SHP-1. LIR-2 mediates tolerogenic DC-induced CD4+ T cell energy in vitro and in vivo.