

## Recombinant Rat IGFBP1/IGFBP-1 protein (His Tag)

Catalog Number: PDER100231

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Rat
<b>Source</b>	E.coli-derived Rat IGFBP1 protein Ala26-Asn272, with an N-terminal His
<b>Mol_Mass</b>	27.1 kDa
<b>Accession</b>	P21743
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 10 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

The insulin-like growth factor binding protein (IGFBP) family consists of six structurally related proteins that bind IGF with high affinity. These proteins share conserved cysteine-rich N- and C-terminal regions that participate in IGF binding. IGFBPs regulate the bioavailability of IGFs and modulate their biological activities, both positively and negatively. Some IGFBPs also have intrinsic bioactivity that is IGF-independent. Post-translational modifications of the IGFBPs, including glycosylation, phosphorylation and proteolysis, influence IGF binding affinities and tissue localization, affecting both the IGF-dependent and independent functions.

### For Research Use Only