
Elabscience®

Elab Fluor[®] Violet 610 Anti-Rat CD45 Antibody[OX-1]

Catalog Number: E-AB-F1227UT

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Rat
Host	Mouse
lsotype	Mouse lgG1, κ
Clone No.	OX-1
Isotype Control	Elab Fluor [®] Violet 610 Mouse IgG1, к Isotype Control[MOPC-21] [Product E-AB- F09792T]
Conjugation	Elab Fluor [®] Violet 610
Conjugation Information	Elab Fluor [®] Violet 610 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 613 nm (e.g., a 615/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.



Staining of normal Rat splenocytes with Elab Fluor[®] Violet 610 Anti-Rat CD45 Antibody[OX-1] (filled gray histogram) or

Elab Fluor[®] Violet 610 Mouse IgG1, κ Isotype Control (empty black histogram). Total viable cells were used for analysis.

Preparation & Storage	
Storage	Keep as concentrated solution.
Shipping	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze. Ice bag
Antigen Information	
Alternate Names	LCA; Ly-5; T200;Leukocyte common antigen;Ptprc;Receptor-type tyrosine-protein phosphatase C
For Research Use Only	

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u> Tel: 1-832-243-6086 Email:techsupport@elabscience.com

Elabscience®

Uniprot ID Gene ID Background P04157

19265

CD45 is a 180-220 kD protein also known as leukocyte common antigen (LCA). It is a protein tyrosine phosphatase with multiple isoforms differing as a result of alternative splicing of the extracellular domain and glycosylation. CD45 is expressed on all hematopoietic cells except erythrocytes and platelets; isoform expression depends on cell type, activation state, and cell maturation. CD45 functions in signal transduction through T and B cell antigen receptors. CD45 has been shown to interact with various proteins including galectin-1, CD2, CD3, and CD4. The OX-1 antibody has been shown to partially inhibit NK cell-mediated lysis of syngeneic tumor cells in vitro.