

APC Anti-Mouse CD27 Antibody[LG.3A10]

Catalog Number: GFH00322E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Mouse
Host	Armenian Hamster
Isotype	Armenian Hamster IgG
Clone No.	LG.3A10
Isotype Control	APC Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09852E]
Conjugation	APC
Conjugation Information	APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide.


Applications

Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of C57BL/6 murine splenocytes cells with FITC Anti-Mouse B220 Antibody and APC Anti-Mouse CD27 Antibody[LG.3A10] (left) or APC Armenian Hamster IgG Isotype Control (right). Total viable cells were used for analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	T14;S152;Tp55;TNFRSF7
Uniprot ID	P41272
Gene ID	21940

For Research Use Only

Background

CD27 is also known as S152 and T14. A member of the tumor necrosis factor receptor (TNFR) superfamily, CD27 is a 45 kD protein expressed on peripheral T cells, memory B cells, NK cells, and thymocyte subset. Through its ligand, CD70, CD27 plays a key role in T cell and B cell interactions. Additionally, ligation of CD27 on naïve T cells may be important in their maturation to effector cells.