



A Reliable Research Partner in Life Science and Medicine

FITC Anti-Mouse TCR γ/δ Antibody[GL3]

Catalog Number: E-AB-F1282UC

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse

Armenian Hamster Host Isotype Armenian Hamster IgG

Clone No. GL3

FITC Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09853C] Isotype Control

Conjugation

Conjugation Information FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical

filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein

protectant.

Applications

Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 µg/10⁶ cells in 100 µL volume].

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names T cell receptor γ/δ **Uniprot ID** Q96E93:O88713 Gene ID 110066;110067

Background T cell receptor (TCR) is a heterodimer consisting of an α and a β chain (TCR α/β) or a γ

> and a δ chain (TCR y/δ). TCR y/δ belongs to the immunoglobulin superfamily, which is involved in the recognition of certain bacterial and tumor antigens bound to MHC class I. γ/δ TCR associates with CD3 and is expressed on a T cell subset found in the thymus, the intestinal epithelium, and the peripheral lymphoid tissues and peritoneum. Most y/δ T cells are CD4-/CD8- although some are CD8+. T cells expressing the y/δ TCR have been shown to play a role in oral tolerance, tumor-associated tolerance, and autoimmune disease. It has been reported that y/\delta T cells also play a principal role in

antigen presentation.

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Web:www.elabscience.com

Email:techsupport@elabscience.com