A Reliable Research Partner in Life Science and Medicine

# Recombinant Human Carbonic Anhydrase 7/CA7 Protein (His Tag)

Catalog Number: PKSH030890

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Species Human

Source E.coli-derived Human Carbonic Anhydrase 7/CA7 protein Met 1-Ala 264, with an C-

terminal His

 Mol\_Mass
 31.0 kDa

 Accession
 P43166

**Bio-activity** Measured by its esterase activity. The activity is > 20 pmoles/min/µg.

#### **Properties**

**Purity** > 96 % as determined by reducing SDS-PAGE.

**Endotoxin** Please contact us for more information.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data



> 96 % as determined by reducing SDS-PAGE.

### Background

Carbonic anhydrase 7; also known as carbonate dehydratase VII; carbonic anhydrase VII; CA-VII and CA7; is a cytoplasm protein which belongs to the alpha-carbonic anhydrase family. Carbonic anhydrases are a large family of zinc metalloenzymes that catalyze the reversible hydration of carbon dioxide. They participate in a variety of biological processes; including respiration; calcification; acid-base balance; bone resorption; and the formation of aqueous humo r; cerebrospinal fluid; saliva; and gastric acid. Carbonic anhydrases show extensive diversity in tissue distribution and in their subcellular localization. CA7 / CA-VII is predominantly expressed in the salivary glands. Alternative splicing in the coding region results in multiple transcript variants encoding different isoforms.

#### For Research Use Only

Fax: 1-832-243-6017