A Reliable Research Partner in Life Science and Medicine

# Recombinant Human FGF-17 Protein (His Tag)

Catalog Number: PKSH032435

Note: Centrifuge before opening to ensure complete recovery of vial contents.

## Description

Species Human

Source HEK293 Cells-derived Human FGF-17 protein Thr 23-Thr 216, with an C-terminal His

 Calculated MW
 22.6 kDa

 Observed MW
 31 kDa

 Accession
 O60258

**Bio-activity** Measured in a cell proliferation assay using Balb/3T3 mouse embryonic fibroblast

cells. The ED<sub>50</sub> for this effect is 2.1 ug/ml.

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### **Background**

Fibroblast Growth Factor 17 (FGF17) is a member of the heparin-binding growth factors family that is prominently expressed in the cerebellum and cortex. Proteins of this family possess broad mitogenic and cell survival activities and they are involved in a variety of biological processes including embryonic development cell growth; morphogenesis; tissue repair; tumor growth; and invasion. FGF17 plays an important role in the regulation of embryonic development and it acts as signaling molecule in the induction and patterning of the embryonic brain. In addition; FGF17 stimulates the proliferation and activation of cells that express FGF receptors.

### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017