



A Reliable Research Partner in Life Science and Medicine

Mouse PDGF-AA Antibody Pair Set

Catalog No. E-KAB-0638 Applications ELISA

Synonyms PDGF-AA

Kit components & Storage

Title	Specifications	Storage
Mouse PDGF-AA Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year. Avoid
		freeze/thaw cycles.
Mouse PDGF-AA Detection Antibody	1 vial, 50 μL	Store at -20°C for one year. Avoid
(Biotin)		freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0638)		
		Maria DDCE AA Cantum Antihada	Mouse PDGF-AA Detection Antibody	
		Mouse PDGF-AA Capture Antibody	(Biotin)	
Immunogen	Immunogen	Recombinant Rat PDGF-AA protien	Recombinant Rat PDGF-AA protien	
Information	Swissprot	P28576		
Product details	Reactivity	Rat	Rat	
	Host	Mouse	Goat	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5 mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%	
		glycerol; pH 7.5	protective protein; 50% glycerol; pH	
			7.5	
	Purify	Protein A or G	Antigen Affinity	
	Specificity	Detects Rat PDGF-AA in ELISAs.		

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com



A Reliable Research Partner in Life Science and Medicine

Applications

Rat PDGF-AA Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Mouse PDGF-AA Capture	
Capture		Antibody	10
			Marie and Marie
EX IO A	1 1000 1 1000	M PDGE AAD	Optical Density
ELISA	1:1000-1:10000	Mouse PDGF-AA Detection	o ·
Detection		Antibody (Biotin)	0.1
			0.1 1 10 100
			Mouse PDGF-AA Concentration (pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Growth factor that plays an essential role in the regulation of embryonic development, cell proliferation, cell migration, survival and chemotaxis. Potent mitogen for cells of mesenchymal origin. Required for normal lung alveolar septum formation during embryogenesis, normal development of the gastrointestinal tract, normal development of Leydig cells and spermatogenesis. Required for normal oligodendrocyte development and normal myelination in the spinal cord and cerebellum. Plays an important role in wound healing. Signaling is modulated by the formation of heterodimers with PDGFB.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com

Tel: 1-832-243-6086 Fax: 1-832-243-6017