

NADK2 Polyclonal Antibody

catalog number: E-AB-18572

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human;Rat
Immunogen	Fusion protein of human NADK2
Host	Rabbit
Isotype	Rabbit IgG
Purification	Antigen affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications

Recommended Dilution


WB	1:500-1:2000
IHC	1:40-1:200

Data




Western blot analysis of A549 and Hela cell lysates using NADK2 Polyclonal Antibody at dilution of 1:1000

Observed-MW: Refer to figures
Calculated-MW: 49 kDa



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using NADK2 Polyclonal Antibody at dilution of 1:40(×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using NADK2 Polyclonal Antibody at dilution of 1:40(×200)

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

This gene encodes a mitochondrial kinase that catalyzes the phosphorylation of NAD to yield NADP. Mutations in this gene result in 2,4-dienoyl-CoA reductase deficiency. Alternative splicing results in multiple transcript variants.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.8