

## Recombinant Human DAPK3/ZIPK Protein (GST Tag)

**Catalog Number:** PKSH030384

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	Baculovirus-Insect Cells-derived Human DAPK3/ZIPK protein Met 1-Arg 454, with an N-terminal GST
<b>Calculated MW</b>	79.0 kDa
<b>Observed MW</b>	70 kDa
<b>Accession</b>	NP_001339.1
<b>Bio-activity</b>	The specific activity was determined to be 5 nmol/min/mg using MBP as substrate.

### Properties

<b>Purity</b>	> 85 % as determined by reducing SDS-PAGE.
<b>Concentration</b>	Subject to label value.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
<b>Shipping</b>	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
<b>Formulation</b>	Supplied as sterile solution of 20mM Tris, 500mM NaCl, 10mM GSH, pH 7.4

### Data



> 85 % as determined by reducing SDS-PAGE.

### Background

Death-associated protein kinase 3, also known as DAP kinase 3, ZIP-kinase, DAPK3 and ZIPK, is a nucleus and cytoplasm protein which belongs to the protein kinase superfamily, CAMK Ser/Thr protein kinase family and DAP kinase subfamily. DAPK3 / ZIPK contains one protein kinase domain. It is a serine/threonine kinase which acts as a positive regulator of apoptosis. It phosphorylates histone H3 on 'Thr-11' at centromeres during mitosis. DAPK3 / ZIPK is a homodimer or forms heterodimers with ATF4. Both interactions require an intact leucine zipper domain and oligomerization is required for full enzymatic activity. It also binds to DAXX and PAWR, possibly in a ternary complex which plays a role in caspase activation. DAPK3 / ZIPK regulates myosin light chain phosphatase through phosphorylation of MYPT1 thereby regulating the assembly of the actin cytoskeleton, cell migration, invasiveness of tumor cells, smooth muscle contraction and neurite outgrowth. It is involved in the formation of promyelocytic leukemia protein nuclear body (PML-NB), one of many subnuclear domains in the eukaryotic cell nucleus, and which is involved in oncogenesis and viral infection.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017