

A Reliable Research Partner in Life Science and Medicine

Recombinant Histone H3 (CT) Monoclonal Antibody

catalog number: AN301708L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

_				40	
D)	es	cr	ıb	tı	on

Reactivity Human;Rat;Mouse;Rice

Immunogen Synthetic peptide corresponding to the C-terminus of human histone H3 protein

 Host
 Rabbit

 Isotype
 IgG, κ


 Clone
 A416

Purification Protein A purified


Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications	Recommended Dilution		
WB	1:1000-1:5000		
IHC	1:100-1:500		
IP	1:50		
IF	1:100-1:500		


Data




Immunohistochemistry of paraffin-embedded Mouse liver using Histone H3 Monoclonal Antibody at dilution of 1:500. Immunohistochemistry of paraffin-embedded Rat skeletal muscle using Histone H3 Monoclonal Antibody at dilution of 1:500.



Immunofluorescent analysis of (100% Ice-cold methanol) Western Blot with Histone H3 Monoclonal Antibody at fixed HeLa cells using anti-Histone H3 Monoclonal Antibody dilution of 1:5000. Lane 1: HeLa, Lane 2: Neuro-2a, Lane 3: at dilution of 1:500. BRL, Lane 4: Rat kidney, Lane 5: Mouse kidney





Calculated-MW:15 kDa

Observed-MW:15 kDa

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com


Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Western Blot with Histone H3 Monoclonal Antibody at dilution of 1:2000. Lane 1: Rice leaves

Observed-MW:15 kDa Calculated-MW:15 kDa Immunohistochemistry of paraffin-embedded Human cerebrum using Histone H3 Monoclonal Antibody at dilution of 1:500.



Immunohistochemistry of paraffin-embedded Human liver using Histone H3 Monoclonal Antibody at dilution of 1:500.

Immunohistochemistry of paraffin-embedded Human testis using Histone H3 Monoclonal Antibody at dilution of 1:500.

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

Histones are a family of basic proteins that form the core of the nucleosome – the fundamental structural unit of chromatin. In a single nucleosome, core histone proteins H2A, H2B, H3, and H4 form an octamer around which the DNA is tightly wrapped. Histone proteins not only serve to compact chromosomal DNA but also play vital roles in the dynamic and long-term regulation of genes by a wide variety of post-translational modifications (PTMs). These PTMs including acetylation, methylation, phosphorylation, and novel acylations directly affect the accessibility of chromatin to transcription factors and other epigenetic regulators, altering genome stability and gene transcription. Among the histone members, histone H3 is the most extensively modified and its amino-terminal tail structure has been shown to be especially critical for eukaryotic cells.

For Research Use Only