


Recombinant Histone H3 (CT) Monoclonal Antibody

catalog number: AN301708L

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description	
Reactivity	Human;Rat;Mouse;Rice
Immunogen	Synthetic peptide corresponding to the C-terminus of human histone H3 protein
Host	Rabbit
Isotype	Rabbit IgG
Clone	C111330
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.
Applications	Recommended Dilution
WB	1:1000-1:5000
IHC	1:100-1:500
IP	1:50
IF	1:100-1:500
Data	




Immunohistochemistry of paraffin-embedded Mouse liver using Histone H3 Monoclonal Antibody at dilution of 1:500.



Immunohistochemistry of paraffin-embedded Rat skeletal muscle using Histone H3 Monoclonal Antibody at dilution of 1:500.





Immunofluorescent analysis of (100% Ice-cold methanol) fixed HeLa cells using anti-Histone H3 Monoclonal Antibody at dilution of 1:500.



Western Blot with Histone H3 Monoclonal Antibody at dilution of 1:5000. Lane 1: HeLa, Lane 2: Neuro-2a, Lane 3: BRL, Lane 4: Rat kidney, Lane 5: Mouse kidney

Observed-MW:15 kDa
Calculated-MW: 15 kDa



For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com


Fax: 1-832-243-6017

Rev. V1.3


Western Blot with Histone H3 Monoclonal Antibody at dilution of 1:2000. Lane 1: Rice leaves

Observed-MW:15 kDa

Calculated-MW: 15 kDa



Immunohistochemistry of paraffin-embedded Human cerebrum using Histone H3 Monoclonal Antibody at dilution of 1:500.



Immunohistochemistry of paraffin-embedded Human liver using Histone H3 Monoclonal Antibody at dilution of 1:500.

Immunohistochemistry of paraffin-embedded Human testis using Histone H3 Monoclonal Antibody at dilution of 1:500.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

Histones are a family of basic proteins that form the core of the nucleosome – the fundamental structural unit of chromatin. In a single nucleosome, core histone proteins H2A, H2B, H3, and H4 form an octamer around which the DNA is tightly wrapped. Histone proteins not only serve to compact chromosomal DNA but also play vital roles in the dynamic and long-term regulation of genes by a wide variety of post-translational modifications (PTMs). These PTMs including acetylation, methylation, phosphorylation, and novel acylations directly affect the accessibility of chromatin to transcription factors and other epigenetic regulators, altering genome stability and gene transcription. Among the histone members, histone H3 is the most extensively modified and its amino-terminal tail structure has been shown to be especially critical for eukaryotic cells.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.3