

## Recombinant Mouse AGO2/Argonaute 2/EIF2C2 Protein (His Tag)

**Catalog Number:** PKSM040551

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Mouse
<b>Source</b>	Baculovirus-Insect Cells-derived Mouse AGO2/Argonaute 2/EIF2C2 protein Met 1-Ala 860, with an N-terminal His
<b>Calculated MW</b>	99.0 kDa
<b>Observed MW</b>	105 kDa
<b>Accession</b>	Q8CJG0
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 80 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile 20mM Tris, 500mM NaCl, 10% Glycerol pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 80 % as determined by reducing SDS-PAGE.

### Background

### For Research Use Only

Argonaute 2 (AGO2), also known as Eukaryotic translation initiation factor 2C2 (EIF2C2), belongs to the Argonaute family, AGO subfamily, which is a component of the RNA-induced silencing complex (RISC) and mediates small interfering RNA (siRNA)-directed mRNA cleavage and microRNA translational suppression. AGO2 protein is the catalytic engine of mammalian RNAi. It contains a PIWI domain that is structurally related to RNases H and possibly shares with them a two-metal-ion catalysis mechanism. Human AGO2 was unable to cleave preformed RNA duplexes and exhibited weaker binding affinity for RNA duplexes compared with the single strand RNA. The enzyme exhibited greater RNase H activity in the presence of Mn<sup>2+</sup> compared with Mg<sup>2+</sup>. Human AGO2 exhibited weaker binding affinities and reduced cleavage activities for antisense RNAs with either a 5'-terminal hydroxyl or abasic nucleotide. In mouse hematopoiesis, AGO2 controls early development of lymphoid and erythroid cells. AGO2 is a highly specialized member of the Argonaute family with an essential nonredundant Slicer-independent function within the mammalian miRNA pathway. AGO2 is a regulator of dFMR1 expression and has an important developmental role for AGO2 in the nervous system and germ line that requires dFMR1 function. In addition, AGO2 is regulated at both the transcriptional and posttranslational level, and also implicate AGO2 and enhanced micro-RNA activity in the tumorigenic progression of breast cancer cell lines.