A Reliable Research Partner in Life Science and Medicine

Recombinant Human Ezrin/EZR Protein

Catalog Number: PKSH033684

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source E.coli-derived Human Ezrin; EZR protein Met1-Leu586

 Mol_Mass
 69.4 kDa

 Accession
 P15311

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin $\leq 1.0 \, \text{EU} \, \text{per} \, \mu \text{g} \, \text{of the protein as determined by the LAL method.}$

Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.


Shipping This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel

packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 µm filtered solution of 10mM HEPES, pH 7.4.

Reconstitution Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Ezrin is expressed in cerebral cortex, basal ganglia, hippocampus, hypophysis, and optic nerve. The N-terminus of ezrin contains a FERM domain which is further subdivided into three subdomains. The C-terminus contain a ERM domain. As a member of the ERM protein family, Ezrin serves as an intermediate between the plasma membrane and the actin cytoskeleton. It plays a key role in cell surface structure adhesion, migration, and organization. Ezrin probably involved in connections of major cytoskeletal structures to the plasma membrane. The N-terminal FERM domain strongly binds sodium-hydrogen exchanger regulatory factor (NHERF) proteins (involving long-range allostery). The C-terminal binds to actin, phosphatidylinositol bis-phosphate (PIP2) and membrane proteins like CD44 and ICAM-2. In epithelial cells, Ezrin is required for the formation of microvilli and membrane ruffles on the apical pole. Along with PLEKHG6, Ezrin is required for normal macropinocytosis.

For Research Use Only

Fax: 1-832-243-6017