## **Elabscience**®

## HSD17B12 Polyclonal Antibody

## catalog number: E-AB-15104

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description   |   |   |
|---|---|---|
| Reactivity  | Human;Mouse;Rat   |   |
| Immunogen   | Recombinant protein of human HSD17B12   |   |
| Host  | Rabbit  |   |
| Is otype  | IgG   |   |
| Purification  | Affinity purification   |   |
| Buffer  | Phosphate buffered sol  | lution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.   |
| Applications  | Recommended Dilution  |   |
| WB  | 1:500-1:2000  |   |
| IHC   | 1:50-1:200  |   |
| Data  |   |   |
| Western Blot analysis of A17<br>HSD17B12 Polyclonal Antibo<br>Calculated-MV | ody at dilution of 1:650  | Immunohistochemistry of paraffin-embedded Human liver<br>cancer using HSD17B12 Polyclonal Antibody at dilution of<br>1:40 |
| Preparation & Storage   |   |   |
| Storage   | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.               |   |
| Shipping  | The product is shipped with ice pack,upon receipt,store it immediately at the |   |
|   | temperature recommended.  |   |
| Background  |   |   |

This gene encodes a very important 17beta-hydroxysteroid dehydrogenase (17beta-HSD) that converts estrone into estradiol in ovarian tissue. This enzyme is also involved in fatty acid elongation. Catalyzes the transformation of estrone (E1) into estradiol (E2), suggesting a central role in estrogen formation. Its strong expression in ovary and mammary gland suggest that it may constitute the major enzyme responsible for the conversion of E1 to E2 in women.