# **PCNA Polyclonal Antibody**

catalog number: E-AB-40489



Note: Centrifuge before opening to ensure complete recovery of vial contents.

	escrip				7.0		
	00		TO T	m	TT.		т
v		ж			ш	w	٠

**Reactivity** Human

Immunogen Synthetic peptide of Human Proliferating cell nuclear antigen

Host Rabbit Isotype IgG


**Purification** Antigen Affinity Purification

**Conjugation** Unconjugated

buffer PBS with 0.05% proclin 300, 1% protective protein and 50% glycerol,pH7.4

**WB** 1:3000-1:7000

## Data



Western Blot analysis of Jurkat, A431 and 293T cells using

PCNA Polyclonal Antibody at dilution of 1:6000.

Observed-MV:35 kDa Calculated-MV:28 kDa

### **Preparation & Storage**

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

#### Background

Proliferating Cell Nuclear Antigen, commonly known as PCNA, is a protein that acts as a processivity factor for DNA polymerase  $\delta$  in eukaryotic cells. This protein is an auxiliary protein of DNA polymerase delta and is involved in the control of eukaryotic DNA replication by increasing the polymerase's processibility during elongation of the leading strand. PCNA induces a robust stimulatory effect on the 3'-5' exonuclease and 3'-phosphodiesterase, but not apurinic-apyrimidinic (AP) endonuclease, APEX2 activities. It has to be loaded onto DNA in order to be able to stimulate APEX2. PCNA protein is highly conserved during evolution; the deduced amino acid sequences of rat and human differ by only 4 of 261 amino acids. PCNA has been used as loading control for proliferating cells.

### For Research Use Only