



A Reliable Research Partner in Life Science and Medicine

Mouse MDC/CCL22 Antibody Pair Set

Catalog No. E-KAB-0317 Applications ELISA

Synonyms CCL22, ABCD-1, DC/B-CK, SCYA22, STCP-1

Kit components & Storage

Title	Specifications	Storage
Mouse MDC/CCL22 Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Mouse MDC/CCL22 Detection Antibody	1 vial, 50 μL	Store at -20℃ for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0317)	
		Mouse MDC/CCL22 Capture	Mouse MDC/CCL22 Detection
		Antibody	Antibody (Biotin)
Immunogen	Immunogen	Recombinant Mouse MDC/CCL22	Recombinant Mouse MDC/CCL22
Information		protein	protein
	Swissprot	O88430	
Product details	Reactivity	Mouse	Mouse
	Host	Rat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Protein A or G	Antigen Affinity
	Specificity	Detects Mouse MDC/CCL22 in ELISAs.	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com





A Reliable Research Partner in Life Science and Medicine

Applications

Mouse MDC/CCL22 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4μg/mL	Mouse MDC/CCL22 Capture	
Capture		Antibody	10 Aug 1
ELISA Detection	1:1000-1:10000	Mouse MDC/CCL22 Detection Antibody (Biotin)	0.01 100 1000 10000 Mouse MDC/CCL22 concentration(pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Macrophage-derived chemokine (MDC), also named stimulated T cell chemotactic protein (STCP-1) and ABCD-1, and now designated as CCL22, is a CC chemokine initially isolated from clones of monocyte-derived macrophages. At the amino acid sequence level, MDC shows less than 35% identity to other CC chemokine family members. Human MDC is expressed in dendritic cells, macrophages and activated monocytes. In addition, MDC expression is detected in thymus, lymph node and appendix tissues. At the amino acid sequence level, mouse and human MDC share 64% identity and 83% similarity.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com