



A Reliable Research Partner in Life Science and Medicine

Human E-Cad Antibody Pair Set

Catalog No. E-KAB-0176 Applications ELISA

Synonyms CDH1, Arc-1, CD324, CDHE, LCAM, UVO, CAM 120/80, Epithelial Cadherin, Uvomorulin

Kit components & Storage

Title	Specifications	Storage
Human E-Cad Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Human E-Cad Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0176)		
		Human E-Cad Capture Antibody	Human E-Cad Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Human E-Cad protein	Recombinant Human E-Cad protein	
Information Swissprot		P12830		
Product details Reactivity		Human	Human	
	Host	Goat	Goat	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%	
		glycerol, pH 7.4	protective protein, 50% glycerol, pH	
P			7.4	
	Purify	Antigen Affinity	Antigen Affinity	
	Specificity	Detects Human E-Cad in ELISAs.	·	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com





A Reliable Research Partner in Life Science and Medicine

Applications

Human E-Cad Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5 -4 μ g/mL	Human E-Cad Capture Antibody	
Capture			Aiss
ELISA Detection	1:1000-1:10000	Human E-Cad Detection Antibody (Biotin)	O.01 O.01 O.01 O.01 Human E-Cad concentration(ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

This gene is a classical cadherin from the cadherin superfamily. The encoded protein is a calcium dependent cell-cell adhesion glycoprotein comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. Mutations in this gene are correlated with gastric, breast, colorectal, thyroid and ovarian cancer. Loss of function is thought to contribute to progression in cancer by increasing proliferation, invasion, and/or metastasis. The ectodomain of this protein mediates bacterial adhesion to mammalian cells and the cytoplasmic domain is required for internalization. Identified transcript variants arise from mutation at consensus splice sites.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com