Elabscience®

p53 Polyclonal Antibody

catalog number: E-AB-93244


Note: Centrifuge before opening to ensure complete recovery of vial contents.

1:50-1:200

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human p53
Host	Rabbit
Is otype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000

Data

IHC




Western blot analysis of extracts of various cell lines using

p53 Polyclonal Antibody at 1:1000 dilution.

Observed-MW:53 kDa

Calculated-MW:23-43 kDa



Western blot analysis of extracts of Rat lung using p53 Polyclonal Antibody at 1:1000 dilution. Observed-MW:53 kDa

60kDa

45kD

Calculated-MW:23-43 kDa

Immunohistochemistry of paraffin-embedded human esophageal cancer using [KO Validated] p53 Polyclonal Antibody at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

before commencing with fire stamming protocol.	
Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.	
The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.	

Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:techsupport@elabscience.com

Elabscience®

This gene encodes a tumor suppressor protein containing transcriptional activation, DNA binding, and oligomerization domains. The encoded protein responds to diverse cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. Mutations in this gene are associated with a variety of human cancers, including hereditary cancers such as Li-Fraumeni syndrome. Alternative splicing of this gene and the use of alternate promoters result in multiple transcript variants and isoforms. Additional isoforms have also been shown to result from the use of alternate translation initiation codons from identical transcript variants (PMIDs: 12032546, 20937277).

For Research Use Only