A Reliable Research Partner in Life Science and Medicine

# Recombinant Rat SCF/c-Kit Ligand Protein (Human Cells, His Tag)

Catalog Number: PKSR030459

Note: Centrifuge before opening to ensure complete recovery of vial contents.

	escrip			7.0	
10)	00	OF	m	177	nn
v	$\mathbf{c}$			U	

**Species** Rat

Source HEK293 Cells-derived Rat SCF/c-Kit Ligand protein Gln26-Ala189, with an C-terminal

His

Calculated MW 19.4 kDa
Observed MW 23-40 kDa
Accession P21581

**Bio-activity** Not validated for activity

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

# Data



> 95 % as determined by reducing SDS-PAGE.

## Background

SCF/C-kit ligand is the ligand of the tyrosine-kinase receptor encoded by the KIT locus. Plays an essential role in the regulation of cell survival and proliferation, hematopoiesis, stem cell maintenance, gametogenesis, mast cell developmen t, migration and function, and in melanogenesis. KITLG/SCF binding can activate several signaling pathways. Promotes phosphorylation of PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase, and subsequent activation of the kinase AKT1. In phase I/II clinical studies administration of the combination of SCF and G-CSF resulted in a two- to threefold increase in cells that express the CD34 antigen compared with G-CSF alone.

#### For Research Use Only