

## Recombinant Human NCAM1 Protein (His Tag)

**Catalog Number:** PKSH032793

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human NCAM1 protein Leu20-Pro603, with an C-terminal His
<b>Calculated MW</b>	65.5 kDa
<b>Observed MW</b>	90-120 kDa
<b>Accession</b>	P13591-3
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Neural cell adhesion molecule 1 (NCAM-1) is a single-pass type I membrane protein, it belongs to a family of membrane-bound glycoproteins that are involved in Ca<sup>2+</sup> independent cell matrix and homophilic or heterophilic cell-cell interactions. NCAM-1 is synthesized as a 761 aa preproprecursor that contains a 19 aa signal sequence, a 722 aa GPI-linked mature region, and a 20 aa C-terminal prosegment. The molecule contains five C-2 type Ig-like domains and two fibronectin type-III domains. NCAM-1 is a cell adhesion molecule involved in neuron-neuron adhesion, neurite fasciculation, outgrowth of neurites, etc. Acting as a receptor for rabies virus, NCAM-1 in the adult brain shows a decline of sialylation relative to earlier developmental periods.

### For Research Use Only