

## Recombinant Human TPBG/5T4 (C-Avi-6His) Biotinylated

**Catalog Number:** PKSH033889

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human TPBG/5T4 protein Ser32-Ser355, with an C-terminal Avi & His
<b>Calculated MW</b>	38 kDa
<b>Observed MW</b>	50-90 kDa
<b>Accession</b>	Q13641
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
<b>Reconstitution</b>	Please refer to the specific buffer information in the printed manual.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Trophoblast glycoprotein, also known as TPBG and 5T4, it belongs to the LRR family of proteins. The C-terminus of LRR N-terminal cap (LRRNT) and LRR 1 are essential for the inhibition of the Wnt signaling pathway. TPBG may function as an inhibitor of Wnt/beta-catenin signaling by indirectly interacting with LRP6 and blocking Wnt3a-dependent LRP6 internalization. It expressed by all types of trophoblasts as early as 9 weeks of development. In adult tissues, the expression is limited to a few epithelial cell types but is found on a variety of carcinoma.

### For Research Use Only