A Reliable Research Partner in Life Science and Medicine

Recombinant Human IL-20RB Protein (Fc Tag)

Catalog Number: PKSH032574

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source HEK293 Cells-derived Human IL-20RB protein Asp30-Ala230, with an C-terminal Fc

Calculated MW49.6 kDaObserved MW60-85 kDaAccessionQ6UXL0

Bio-activity Not validated for activity

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Concentration Subject to label value.

Endotoxin $< 1.0 \text{ EU per } \mu\text{g}$ of the protein as determined by the LAL method.

Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.


This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel

packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Data

Shipping



> 90 % as determined by reducing SDS-PAGE.

Background

Interleukin-20 receptor subunit beta (IL20RB) is a single-pass type I membrane protein and belongs to the type II cytokine receptor family. It contains 2 fibronectin type-III domains. There are two kinds of type II cytokine receptors: cytokine receptors that bind type I and type II interferons; cytokine receptors that bind members of the interleukin-10 family (interleukin-10, interleukin-20 and interleukin-22). Type II cytokine receptors are similar to type I cytokine receptors except they do not possess the signature sequence WSXWS that is characteristic of type I receptors. They are expressed on the surface of certain cells, which bind and respond to a select group of cytokines. These receptors are related predominantly by sequence similarities in their extracellular portions that are composed of tandem Ig-like domain s. The intracellular domain of type II cytokine receptors is typically associated with a tyrosine kinase belonging to the Janus kinase (JAK) family