

A Reliable Research Partner in Life Science and Medicine

Elab Fluor[®] 488 Anti-Human TCRVγ9 Antibody[B3]

Catalog Number: AN00357L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human
Host Mouse
Isotype Mouse IgG1

Clone No. B3

Isotype Control Elab Fluor® 488 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792L]

Conjugation Elab Fluor® 488

Conjugation Information Elab Fluor[®] 488 is designed to be excited by the Blue laser (488 nm) and detected using

an optical filter centered near 520 nm (e.g., a 525/40 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of normal human peripheral blood cells with APC

Anti-Human CD3 Antibody and Elab Fluor[®] 488 Anti-Human TCR Vγ9 Antibody[B3] (left) or Elab Fluor[®] 488 Mouse IgG1, κ Isotype Control (right). Cells in the lymphocytes gate were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names T cell receptor Vy9;TCR Vy2;TRGV9;TCRGV2S1

 Uniprot ID
 Q99603

 Gene ID
 6965

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Background

Elabscience®

The Vγ9 TCR is a variant of the TCR γ chain expressed on a subset of γ/δ T cells. Vγ9Vδ 2 T lymphocytes, a major y/δ T cell subset in humans, recognize phosphoantigens, certain tumor cells, and cells treated with aminobisphosphonates. This cell population displays cytolytic activity against various tumor cells. The γ/δ TCR is a heterodimeric TCR complex composed of covalently bound γ and δ chains involved in antigen recognition and the non-covalently associated monomorphic proteins CD3 δ , γ , ϵ , and ζ chains.

Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Email:techsupport@elabscience.com Web:www.elabscience.com