A Reliable Research Partner in Life Science and Medicine

Myc-Tag Monoclonal Antibody

catalog number: E-AB-48023

Note: Centrifuge before opening to ensure complete recovery of vial contents.

\mathbf{r}					
D	es	cr	П	IIC	or

Reactivity All

Immunogen Synthetic peptide corresponding to Myc tag conjugated to keyhole limpet

haemocyanin.

HostMouseIsotypeIgG2aClone3E9

Purification Protein A/G Purification

Conjugation Unconjugated


Buffer PBS, pH 7.2. Contains 0.05% Proclin300.

Applications Recommended Dilution

WB 1:2000-4000 FCM 0.2μg

Data





Western blotting with anti-MYC Mouse monoclonal antibody at dilution of 1:1000.Lane 1: MYC tag transfected HEK 293 cell lysates, lane 2: HEK 293 whole cell lysate

Observed-MW:28 kDa Calculated-MW:28 kDa 1x10^6 CHO cells Transfected with a MYC plasmid were stained with 0.2ug Anti-MYC tag mAb(E-AB-48023) and AF647 conjugated Goat Anti-mouse IgG(H+L);lsotype Control stained with 0.2ug mouse IgG and AF647 conjugated Goat Anti-mouse IgG (H+L)

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack,upon receipt, store it immediately at the

temperature recommended.

Background

The MYC tag, derived from the c-MYC protein, is a popular epitope tag for detecting the expression of recombinant proteins in yeast, bacteria, insect, and mammalian cell systems. It can be used for affinity chromatography, then used to separate recombinant, overexpressed protein from wild type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits.