

PE/Cyanine5 Anti-Rat CD172a(SIRPα) Antibody[OX-41]

Catalog Number: AN00652G

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Rat
Host	Mouse
Isotype	Mouse IgG2a, κ
Clone No.	OX-41
Isotype Control	PE/Cyanine5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792G]
Conjugation	PE/Cyanine 5
Conjugation Information	PE/Cyanine 5 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 670 nm (e.g., a 690/50 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of C57BL/6 rat bone marrow cells with PE/Cyanine 5 Anti-Rat CD172a(SIRPα) Antibody[OX-41] (filled gray histogram) or PE/Cyanine5 Mouse IgG2a, κ Isotype Control (empty black histogram). Total viable cells were used for analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	SIRPα, PTPNS1, BIT, MFR, p84, CD172 antigen-like family member A, AN00652
Uniprot ID	P97710

For Research Use Only

Gene ID

19261

Background

Rat CD172a, also known as a signal-regulatory protein 伪 (SIRP伪), is a 75-110 kD transmembrane glycoprotein involved in the receptor tyrosine kinase coupled signaling pathway. It belongs to the Ig superfamily primarily expressed on monocytes/macrophages, granulocytes, dendritic cells and neurons. CD172a serves as a substrate of activated receptor tyrosine kinases (RTKs). The interaction of the CD172a intracellular domain with SHP-1 and SHP-2 displays negative signaling in the regulation of leukocyte adhesion and transmigration, T cell activation, macrophage fusion and phagocytosis. CD47 (IAP) is an extracellular ligand for CD172a.