

PE/Cyanine5 Anti-Rat CD172a(SIRPa) Antibody[OX-41]

Catalog Number: AN00652G

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Rat
Host	Mouse
Isotype	Mouse IgG2a, κ
Clone No.	OX-41
Isotype Control	PE/Cyanine5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792G]
Conjugation	PE/Cyanine 5
Conjugation Information	PE/Cyanine 5 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 670 nm (e.g., a 690/50 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	---

Data



Staining of C57BL/6 rat bone marrow cells with PE/Cyanine 5 Anti-Rat CD172a(SIRPa) Antibody[OX-41] (filled gray histogram) or PE/Cyanine5 Mouse IgG2a, κ Isotype Control (empty black histogram). Total viable cells were used for analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protect from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	SIRPa, PTPNS1, BIT, MFR, p84, CD172 antigen-like family member A;AN00652
Uniprot ID	P97710

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.2

Gene ID	19261
Background	<p>Rat CD172a, also known as a signal-regulatory protein 伪 (SIRP伪), is a 75-110 kD transmembrane glycoprotein involved in the receptor tyrosine kinase coupled signaling pathway. It belongs to the Ig superfamily primarily expressed on monocytes/macrophages, granulocytes, dendritic cells and neurons. CD172a serves as a substrate of activated receptor tyrosine kinases (RTKs). The interaction of the CD172a intracellular domain with SHP-1 and SHP-2 displays negative signaling in the regulation of leukocyte adhesion and transmigration, T cell activation, macrophage fusion and phagocytosis. CD47 (IAP) is an extracellular ligand for CD172a.</p>

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.2