Recombinant Human NME1/NDKA Protein (His Tag)

Catalog Number: PKSH032830



Note: Centrifuge before opening to ensure complete recovery of vial contents.

		crip				
	00	CI	РΤ	n	tπ	Λn
JU.	\mathbf{c}	v.		w	ш	UП

 Species
 Human

 Mol_Mass
 19.3 kDa

 Accession
 P15531

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin $\leq 1.0 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$ **Storage** Storage Stor

Shipping This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel


packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 1mM DTT, 10% Glycerol,

pH 7.5.

Reconstitution Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Nucleoside-Diphosphate Kinases (NDKs) are enzymes that catalyze the exchange of phosphate groups between different nucleoside diphosphates. NDKs Possesse nucleoside-diphosphate kinase, serine/threonine-specific protein kinase, geranyl and farnesyl pyrophosphate kinase, histidine protein kinase and 3-5 exonuclease activities. NDKs involved in cell proliferation, differentiation and development, signal transduction, G protein-coupled receptor endocytosis, and gene expression and required for neural development including neural patterning and cell fate determination. Prokaryotic NDK forms a functional homotetramer. There are two isoforms of NDK in humans: NDK-A and NDK-B. Both have very similar structure, and can combine in any proportion to form functional NDK hexamers.

For Research Use Only