

A Reliable Research Partner in Life Science and Medicine

Recombinant CD74 Monoclonal Antibody

catalog number: AN301823L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human;

Immunogen Recombinant human CD74 fragment

HostRabbitIsotypeIgG, κCloneA535


Purification Protein A purified

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution

WB 1:500-1:2000 IHC 1:2000-1:5000 FCM 1:50


Data



Western Blot with CD74 Monoclonal Antibody at dilution of Immunohistochemistry of paraffin-embedded Human spleen 1:2000. Lane 1: Jurkat (Negative control), Lane 2: Raji, Lane using CD74 Monoclonal Antibody at dilution of 1:5000.

3: Ramos


Observed-MW:34 kDa Calculated-MW:34 kDa



Immunohistochemistry of paraffin-embedded Human tonsil using CD74 Monoclonal Antibody at dilution of 1:2000.

Flow cytometric analysis of human CD74 expression on Ramos cells. Cells were stained with purified anti-Human CD74, then a Alexa Fluor 488-conjugated second step antibody. The histogram were derived from events with the forward and side light-scatter characteristics of intact cells.

Rev. V1.1



For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Elabscience®

Flow cytometric analysis of human CD74 expression on Jurkat(Negative cell line) cells. Cells were stained with purified anti-Human CD74, then a Alexa Fluor 488-conjugated second step antibody. The histogram were derived from events with the forward and side light-scatter characteristics of intact cells.

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

CD74, also known as the MHC Class II-associated invariant chain (Ii), is a type II transmembrane glycoprotein that plays a critical role in the antigen presentation process as a chaperone of MHC Class II proteins. It is expressed at high levels on B cells and to a lesser extent on numerous antigen presenting cell (APC) types including dendritic cell s, Langerhans cells, monocytes, and macrophages as well as non-traditional APCs such as epithelial cells. CD74 was initially identified for its ability to regulate the folding and intracellular trafficking of newly synthesized MHC Class II molecules. Following expression, CD74 self-assembles as a trimer that serves as a scaffold for the assembly of MHC Class II molecules. Through this interaction, CD74 blocks the peptide binding cleft of MHC Class II molecules and prevents their premature association with endogenous polypeptides. Binding to CD74 also facilitates the translocation of MHC Class II molecules from the endoplasmic reticulum to the endocytic compartments during antigen presentation.