

A Reliable Research Partner in Life Science and Medicine

PerCP Anti-Mouse CD8a Antibody[53-6.7]

Catalog Number: E-AB-F1104F

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Rat Host

Isotype Rat IgG2a, ĸ Clone No. 53-6.7

PerCP Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832F] Isotype Control

PerCP Conjugation

Conjugation Information PerCP is designed to be excited by the blue laser (488 nm) and detected using an optical

filter centered near 675 nm (e.g., a 690/50 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



C57BL/6 murine splenocytes are stained with PerCP Anti-Mouse CD8a Antibody (filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.

Preparation & Storage

Keep as concentrated solution. Storage

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD8A;MAL;T-cell surface glycoprotein CD8 alpha chain;T-lymphocyte differentiation

antigen T8/Leu-2

Uniprot ID P01731 Gene ID 12525

For Research Use Only

Fax: 1-832-243-6017 Tel: 1-832-243-6086 Toll-free: 1-888-852-8623 Web:www.elabscience.com

Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Background

CD8, also known as Lyt-2, Ly-2, or T8, consists of disulfide-linked α and β chains that form the α (CD8a)/ β (CD8b) heterodimer and α / α homodimer. CD8a is a 34 kD protein that belongs to the immunoglobulin family. The CD8 α / β heterodimer is expressed on the surface of most thymocytes and a subset of mature TCR α / β T cells. CD8 expression on mature T cells is non-overlapping with CD4. The CD8 α / α homodimer is expressed on a subset of γ / δ TCR-bearing T cells, NK cells, intestinal intraepithelial lymphocytes, and lymphoid dendritic cells. CD8 is an antigen co-receptor on T cells that interacts with MHC class I on antigen-presenting cells or epithelial cells. CD8 promotes T cell activation through its association with the TCR complex and protein tyrosine kinase lck.

Fax: 1-832-243-6017