

A Reliable Research Partner in Life Science and Medicine

# PE/Cyanine 5 Anti-Rat CD44H Antibody[OX-49]

Catalog Number: E-AB-F1225UG

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

Reactivity Rat
Host Mouse

**Isotype** Mouse IgG2a, κ

Clone No. OX-49

Isotype Control PE/Cyanine5 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09803G]

Conjugation PE/Cyanine 5

Conjugation Information PE/Cyanine5 is designed to be excited by the Blue (488 nm), Green (532 nm) and

yellow-green (561 nm) lasers and detected using an optical filter centered near 670 nm

(e.g., a 690/50 nm bandpass filter).

**Storage Buffer** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

#### Applications Recommended usage


FCM Each lot of this antibody is quality control tested by flow cytometric analysis. Please

check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the

reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu g/10^6$  cells

in 100 µL volume].

#### Data



Rat splenocytes are stained with PE/Cyanine5 Anti-Rat CD44H Antibody (filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.

#### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

#### **Antigen Information**

Alternate Names CD44sH-CAM;CD44H;Pgp-1

 Uniprot ID
 P26051

 Gene ID
 12505

# For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

#### Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

## **Background**

CD44 is an 80-95 kD cell surface glycoprotein. It is expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. It is up-regulated when T cells and B cells are activated. It was reported that CD44 is a valuable marker for memory T cells. CD44 is an adhesion molecule involved in leukocyte adhesion and homing to lymphoid organs. The OX-49 antibody reacts with CD44H (known as CD44s) expressed on most leukocytes, except for a subset of B lymphocytes. The epitope recognized by OX-49 antibody has been mapped to a region on both the standard, CD44s, and the splice variant, CD44v, isoforms of CD44. However it was reported that OX-49 antibody cannot detect the CD44V isoform, possibly due to conformational changes in the epitope.

Fax: 1-832-243-6017