## **GZMM** Polyclonal Antibody

catalog number: E-AB-53442



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description                       |  |   |  |
|-----------------------------------|--|---|--|
| Description                       |  |   |  |
| Reactivity                        | Human;Mouse;Rat  |   |  |
| Immunogen                         | Synthetic peptide of human GZMM  |   |  |
| Host                              | Rabbit   |   |  |
| Isotype                           | IgG  |   |  |
| Purification                      | Antigen affinity purification  |   |  |
| Conjugation                       | Unconjugated   |   |  |
| buffer                            | Phosphate buffered solu  | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.  |  |
| Applications                      | Recommended Dilution   |   |  |
| WB                                | 1:500-1:2000   | 1:500-1:2000  |  |
| IHC                               | 1:20-1:100   |   |  |
| Data                              |  |   |  |
| Polyclonal Antibod<br>Observed-MV | AW264.7 cell using GZMM<br>y at dilution of 1:400<br><b>Refer to figures</b><br>- <b>MV:28 kDa</b>       | Immunohistochemistry of paraffin-embedded Human brain<br>tissue using GZMM Polyclonal Antibody at dilution of<br>1:25(×200) |  |
| Preparation & Storage             |  |   |  |
| Storage                           | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.  |   |  |
| Shipping                          | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |   |  |

## Background

GZMM (Granzyme M) is a Protein Coding gene. Diseases associated with GZMM include Adrenal Insufficiency, Congenital, With 46Xy Sex Reversal, Partial Or Complete and Congenital Adrenal Insufficiency. Among its related pathways are Innate Immune System and Granzyme Pathway. GO annotations related to this gene include serine-type endopeptidase activity and endopeptidase activity. An important paralog of this gene is CFD.Human natural killer (NK) cells and activated lymphocytes express and store a distinct subset of neutral serine proteases together with proteoglycans and other immune effector molecules in large cytoplasmic granules. These serine proteases are collectively termed granzymes and include 4 distinct gene products: granzyme A, granzyme B, granzyme H, and the protein encoded by this gene, granzyme M. Two transcript variants encoding different isoforms have been found for this gene.

## For Research Use Only