


Recombinant BAP29 Monoclonal Antibody

catalog number: **AN301714L**


Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description	
Reactivity	Human
Immunogen	Recombinant human BAP29 fragment
Host	Rabbit
Isotype	Rabbit IgG
Clone	C311336
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.
Applications	Recommended Dilution
WB	1:1000-1:5000
IHC	1:200-1:1000
IF	1:50
IP	1:50-1:100
Data	




Western Blot with BAP29 Monoclonal Antibody at dilution of 1:5000. Lane 1: Jurkat, Lane 2: U87-MG, Lane 3: SW480


Observed-MW:27 kDa
Calculated-MW:27 kDa




Immunohistochemistry of paraffin-embedded Human testis using BAP29 Monoclonal Antibody at dilution of 1:1000.



Immunohistochemistry of paraffin-embedded Human cerebrum using BAP29 Monoclonal Antibody at dilution of 1:1000.



Immunofluorescent analysis of (100% methanol) fixed HeLa cells using anti-BAP29 Monoclonal Antibody at dilution of 1:50.



For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.2

Immunoprecipitation analysis using anti-BAP29
Monoclonal Antibody. Western blot was performed from
the immunoprecipitate using BAP29 Monoclonal
Antibody at a dilution of 1:100. Lane 1: 5% Input, Lane 2:
BAP29 Monoclonal Antibody, Lane 3: Rabbit monoclonal
IgG Isotype

Observed-MW:27 kDa

Calculated-MW:27 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

This protein may play a role in anterograde transport of membrane proteins from the endoplasmic reticulum to the Golgi and be involved in CASP8-mediated apoptosis.