

Elab Fluor® 488 Anti-Human CD193/CCR3 Antibody[5E8]

Catalog Number: E-AB-F1303L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Human
Host	Mouse
Isotype	Mouse IgG2b, κ
Clone No.	5E8
Isotype Control	Elab Fluor® 488 Mouse IgG2b, κ Isotype Control[MPC-11] [Product E-AB-F09812L]
Conjugation	Elab Fluor® 488
Conjugation Information	Elab Fluor® 488 is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 520 nm (e.g., a 525/40 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.


Applications

Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Human peripheral blood granulocytes are stained with APC Anti-Human CD16 Antibody and Elab Fluor® 488 Anti-Human CD193/CCR3 Antibody (Left). Granulocytes are stained with APC Anti-Human CD16 Antibody and Elab Fluor® 488 Mouse IgG2b,κ Isotype Control (Right).

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	MIP1-alpha receptor like-2/eotaxin receptor;C-C chemokine receptor type 3;CC CKR3; CCR;CCR3
Uniprot ID	P51677

For Research Use Only

Gene ID

1232

Background

CD193, also known as CC-chemokine receptor 3 (CCR3), CC CKR3, MIP1-alpha receptor like-2, and eotaxin receptor, is a member of the G protein-coupled seven transmembrane receptors family. It binds to the CC chemokines eotaxin, eotaxin-2, and eotaxin-3 with high affinity. CCR3 has also been reported to bind RANTES, MCP-3, and MCP-4 with low affinity. CCR3 receptor is expressed on human eosinophils, basophils, mast cells, mononuclear phagocytes, platelets, CD34+ hematopoietic progenitor cells, Th2-like lymphocytes, and keratinocytes. CCR3 is thought to play a role in allergic diseases such as bronchial asthma and allergic rhinitis. CCR3 is a co-receptor for HIV-1 and HIV-2, and the binding of eotaxin with CCR3 has been shown to inhibit HIV infection in some cell types.