Recombinant Human CD72/Lyb-2 Protein (Trx&His)

Catalog Number: PKSH032112



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| - | | | | | |
|------|----|--------------|---|-----|----|
| - 10 | 00 | cri | m | 17 | ٦m |
| u | | \mathbf{u} | | 7 U | |

 Species
 Human

 Mol_Mass
 30.6 kDa

 Accession
 P21854

Bio-activity Not validated for activity

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs. **Formulation**Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

B-Cell Differentiation Antigen CD72 (CD72) is a single-pass type II membrane protein. CD72 exists as a disulfide-linked homodimer and contains one C-type lectin domain. CD72 is expressed on B lineage cells, NK cells, monocytes, dendritic cells, and mast cells. CD72 is a ligand for CD5. CD72 associates with CD5, interacts with PTPN6/SHP-1 and plays a role in B-cell proliferation and differentiation. CD72 associates with CD79A in the B cell antigen receptor (BCR) complex following antigen stimulation and dampens BCR signaling through interactions with the phosphatase SHP-1.

For Research Use Only