

A Reliable Research Partner in Life Science and Medicine

## Elab Fluor® 488 Anti-Mouse CD86 Antibody[GL-1]

Catalog Number: E-AB-F0994UL

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Mouse Rat Host

Isotype Rat IgG2a, ĸ

Clone No. GL-1

Isotype Control Elab Fluor® 488 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833L]

Conjugation Elab Fluor®488

**Conjugation Information** Elab Fluor® 488 is designed to be excited by the Blue laser (488 nm) and detected using

an optical filter centered near 520 nm (e.g., a 525/40 nm bandpass filter).


Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. Storage Buffer

#### **Applications** Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μg/10<sup>6</sup> cells in 100 µL volume].

#### Data



C57BL/6 murine splenocytes are stained with Elab Fluor® 488 Anti-Mouse CD86 Antibody (filled gray histogram).

Unstained splenocytes (empty black histogram) are used as control.

#### **Preparation & Storage**

Keep as concentrated solution. **Storage** 

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

**Shipping** Ice bag

#### **Antigen Information**

**Alternate Names** Activation B7-2 antigen; Cd86; ETC-1; Early T-cell costimulatory molecule 1; T-lymphocyte

activation antigen CD86

**Uniprot ID** P42082 Gene ID 12524

### For Research Use Only

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Email:techsupport@elabscience.com

Web:www.elabscience.com

# Elabscience®

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

#### **Background**

CD86 is an 80 kD immunoglobulin superfamily member also known as B7-2, B70, and Ly-58. CD86 is expressed on activated B and T cells, macrophages, dendritic cells, and astrocytes. CD86, along with CD80, is a ligand of CD28 and CD152 (CTLA-4). CD86 is expressed earlier in the immune response than CD80. CD86 has also been shown to be involved in immunoglobulin class-switching and triggering of NK cell-mediated cytotoxicity. CD86 binds to CD28 to transduce co-stimulatory signals for T cell activation, proliferation, and cytokine production. CD86 can also bind to CD152, also known as CTLA-4, to deliver an inhibitory signal to T cells.

Fax: 1-832-243-6017