

A Reliable Research Partner in Life Science and Medicine

FITC Anti-Human CD18 Antibody[TS1/18.1.2.11]

Catalog Number: E-AB-F1057C

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Host Mouse

Isotype Mouse IgG1, κ **Clone No.** TS1/18.1.2.11

Isotype Control FITC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792C]

Conjugation FITC

Conjugation Information FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical

filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Human peripheral blood lymphocytes are stained with FITC Anti-Human CD18 Antibody (filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD18;Cell surface adhesion glycoproteins LFA-1/CR3/p150+95 subunit beta;

Complement receptor C3 subunit beta;Integrin beta-2;Itgb2

 Uniprot ID
 P05107

 Gene ID
 3689

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Background

Elabscience®

CD18 is a 90-95 kD type I transmembrane protein also known as integrin $\beta2$ subunit, LFA-1 β subunit, and $\beta2$ integrin. CD18 non-covalently associates with CD11a, CD11b or CD11c. CD18 is expressed on all leukocytes. CD18 and associated α chains function in adhesion and signaling in hematopoietic cells.

Toll-free: 1-888-852-8623 Web:www.elabscience.com Tel: 1-832-243-6086 Email:techsupport@elabscience.com Fax: 1-832-243-6017