Recombinant Human IL-1RAcP/IL1R3 Protein (Fc & His Tag)

Catalog Number: PKSH032559



| Description | | | |
|----------------|--|--|--|
| Species | Human | | |
| Mol_Mass | 66.7 kDa | | |
| Accession | Q9NPH3-2 | | |
| Bio-activity | Not validated for activity | | |
| Properties | | | |
| Purity | > 95 % as determined by reducing SDS-PAGE. | | |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. | | |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 | | |
| | °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of | | |
| | reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. | | |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. | | |
| Formulation | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. | | |
| | Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants | | |
| | before lyophilization. | | |
| | Please refer to the specific buffer information in the printed manual. | | |
| Reconstitution | Please refer to the printed manual for detailed information. | | |
| Data | | | |

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| kDa | MK |
|-----------|--------------|
| 120 90 | |
| 90 60 | - abbellette |
| 40 | - |
| 30 | - cience |
| 30 20 | Ela |
| 20 | hand |
| | |

> 95 % as determined by reducing SDS-PAGE.

Background

Interleukin-1 Receptor Accessory Protein (IL-1RAcP) is a member of the interleukin-1 receptor family. It contains three I g-like C2-type domains in the extracellular region and a long cytoplasmic domain implicated in signal transduction. IL-1RAcP acts as a non-ligand binding accessory component of the receptors for IL1α; IL1βand IL33. IL-1RAcP mediates interleukin-1-dependent activation of NF-kappa-B. It is part of the membrane-bound form of the IL-1 receptor. IL-1 RAcP takes part in the Signaling ways by the formation of a ternary complex containing IL1R1; TOLLIP; MYD88; and IRAK1 or IRAK2. In addition; IL-1RAcP modulates the response to interleukins by associating with soluble IL1R1 and enhancing interleukin-binding to the decoy receptor.

For Research Use Only