Recombinant Rat Ninjurin-1/NINJ1 Protein (Fc Tag)

Catalog Number: PKSR030174



Note: Centrifuge before opening to ensure complete recovery of vial contents.

		- 40	
Desc	\mathbf{r}	$\mathbf{n}\mathbf{T}\mathbf{I}$	\mathbf{on}
DUSU	ш		ULL

Synonyms NINJ1
Species Rat

Expression Host

Sequence

Met 1-Pro 79

Accession

P70617

Calculated Molecular Weight

Observed molecular weight

Tag

N-hFc

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Ninjurin-1, also known as NINJ1, is a member of the Ninjurin family of transmembrane (TM) proteins. It is expressed in CD19(+) CD10(+) B-cell progenitor cells and higher levels in B-lineage acute lymphoblastic leukemia cells. Ninjurin-1 is expressed also in a number of other adult and embryonic tissues, predominantly in epithelial cells. Its expression is upregulated after axotomy in neurons and in Schwann cells surrounding the distal nerve segment. Upregulated expression of ninjurin-1 has been identified as a marker of minimal residual disease in B-lineage acute lymphoblastic leukemia. It mediates homophilic adhesion, and promotes neurite extension of dorsal root ganglion neurons in vitro. Ninjurin-1 has been found to show a high expression level in the liver tissue of patients with hepatocellular carcinoma, and this seems to be associated with cases of cirrhosis and chronic viral hepatitis. It has been reported that NINJURIN increases p21

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

Recombinant Rat Ninjurin-1/NINJ1 Protein (Fc Tag)

Catalog Number:PKSR030174



expression and induces cellular senescence in human hepatoma cells.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com