## Biotin Anti-Human CD11a Antibody[TS1/22.1.1.13]

## Catalog Number: E-AB-F1055B

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Host	Mouse
lsotype	Mouse lgG1, κ
Clone No.	TS1/22.1.1.13
Isotype Control	Biotin Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09793B]
Conjugation	Biotin
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. For flow
	cytometric staining, the suggested use of this reagent is $\leq 1.0 \ \mu$ g per 10 <sup>6</sup> cells in 100 $\mu$ L volume or 100 $\mu$ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Preparation & Storage	
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	CD11 antigen-like family member A;CD11a;Integrin alpha-L;Itgal;LFA-1A;Ly-15;
	Lymphocyte antigen 15
Uniprot ID	P20701
Gene ID	3683
Background	CD11a is a 170-180 kD type I transmembrane glycoprotein also known as LFA-1 $lpha$
	chain and integrin $\alpha L$ subunit. CD11a non-covalently associates with integrin $\beta 2$ (CD1
	8) to form LFA-1. It is expressed on all leukocytes, including B and T lymphocytes,
	monocytes, macrophages, neutrophils, basophils and eosinophils. It is absent on non-
	hematopoietic tissues and platelets. CD11a plays a central role in leukocyte cell-cell
	interactions and is important in lymphocyte costimulation. CD11a/CD18 binds to ICAM-
	1 (CD54), ICAM-2 (CD102), and ICAM-3 (CD50).

## For Research Use Only