

A Reliable Research Partner in Life Science and Medicine

ICAM-1/CD54 Monoclonal Antibody

catalog number: AN200003P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human ICAM-1 protein


HostMouseIsotypeIgG1CloneA847PurificationProtein A


Buffer 0.2 µm filtered solution in PBS

Applications Recommended Dilution


ICC/IF 1:20-1:100 FCM 1:25-1:100

Data





Flow cytometric analysis of Human ICAM1 (CD54) in HUVEC cells. HUVEC cells (Human umbilical vein endothelial cells) were either treated with TNF-α (50ng/mL, 18 hours) (right panel) or untreated (left panel). The cells were harvested and stained with purified human ICAM-1, then a FITC-conjugated second step antibody.



Immunofluorescence staining of Human ICAM1 in MSC cells. Cells were fixed with 4% PFA,blocked with 10% serum, and incubated with Mouse anti-Human ICAM1 Monoclonal Antibody (1:60) at 37°C 1 hour. Then cells were stained with the Alexa Fluor® 488-conjugated Goat Antimouse IgG secondary antibody (green) and counterstained with DAPI for nuclear staining (blue). Positive staining was localized to cell membrane.

Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

Background

ICAM-1 is a member of the immunoglobulin superfamily whose expression is upregulated on leukocytes, epithelial cells and resting endothelial cells in response to inflammatory signals. ICAM-1 binds the leukocyte integrins LFA-1 and Mac-1.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com