

PerCP/Cyanine5.5 Anti-Rat CD3 Antibody[G4.18]

Catalog Number: E-AB-F1228UJ

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|--------------------------------|--|
| Reactivity | Rat |
| Host | Mouse |
| Isotype | Mouse IgG3, κ |
| Clone No. | G4.18 |
| Isotype Control | PerCP/Cyanine5.5 Mouse IgG3, κ Isotype Control[A112-3] [Product E-AB-F09753J] |
| Conjugation | PerCP/Cyanine 5.5 |
| Conjugation Information | PerCP/Cyanine5.5 is designed to be excited by the blue laser (488 nm) and detected using an optical filter centered near 675 nm (e.g., a 690/50 nm bandpass filter). |
| Storage Buffer | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant. |

Applications

| Applications | Recommended usage |
|--------------|--|
| FCM | Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 µg/10 ⁶ cells in 100 µL volume]. |

Data



Rat splenocytes are stained with PerCP/Cyanine5.5 Anti-Rat CD3 Antibody (filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.

Preparation & Storage

| | |
|-----------------|---|
| Storage | Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze. |
| Shipping | Ice bag |

Antigen Information

| | |
|------------------------|--|
| Alternate Names | T3and ζ chainsγc;CD3;CD3 Complex;T-cell surface glycoprotein CD3 δ |
| Uniprot ID | P19377;Q64159;D4A5M2 |
| Gene ID | 25710,300678,315609,25300 |

For Research Use Only

Background

CD3 is a complex composed of δ , γ , ϵ , and ζ chains. They are 20-25 kD members of the immunoglobulin superfamily and associated with the T cell receptor (TCR). CD3 is expressed on thymocytes, peripheral T cells, some NK-T cells, and dendritic epidermal T cells. CD3 is involved in antigen recognition, signal transduction, and T cell activation.

For Research Use Only