

Purified Anti-Human CD279/PD-1 Antibody[J116]

catalog number: E-AB-F1129A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Immunogen	Recombinant Human CD279 protein
Host	Mouse
Isotype	Mouse IgG1, κ
Clone	J116
Purification	>98%, Protein A/G purified
Buffer	PBS, pH 7.2. Contains 0.05% proclin 300.

Applications

Recommended Dilution

FCM	2 μ g/mL (1×10^5 - 5×10^5 cells)
------------	---

Data



Molt-4 were activated for 4 hour with MIX, then stained with 0.2 μ g Purified Anti-Human CD279 Antibody[J116] (Right) and 0.2 μ g Mouse IgG1, κ Isotype Control(Left), followed by APC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

CD279, also known as programmed death-1 (PD-1), is a 50-55 kD glycoprotein belonging to the CD28 family of the Ig superfamily. PD-1 is expressed on activated splenic T and B cells and thymocytes. It is induced on activated myeloid cells as well. PD-1 is involved in lymphocyte clonal selection and peripheral tolerance through binding its ligands, B7-H1 (PD-L1) and B7-DC (PD-L2). It has been reported that PD-1 and PD-L1 interactions are critical to positive selection and play a role in shaping the T cell repertoire. PD-L1 negative costimulation is essential for prolonged survival of intratesticular islet allografts.

For Research Use Only