

**FGFR-3/CD333/FGFR3, Human, Recombinant****Cat. No. : GPCK139****产品信息**

物种	Human
表达宿主	Human Cells
序列信息	Glu23-Gly375
检索号	P22607
标签	C-6His
分子量	39 kDa
有效期	12 months

**产品特性**

内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
复溶	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**背景介绍**网站: [www.procell.com.cn](http://www.procell.com.cn)

电话: 400-999-2100

邮箱: [techsupport@procell.com.cn](mailto:techsupport@procell.com.cn)

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



Rev. V1.3

Fibroblast Growth Factors (FGFs) comprise a family of at least eighteen structurally related Proteins that are involved in a multitude of physiological and pathological cellular processes, including cell growth, differentiation, angiogenesis, wound healing and tumorigenesis. The biological activities of the FGFs are mediated by a family of type I transmembrane tyrosine kinases which undergo dimerization and autophosphorylation after Ligand binding. Four distinct genes encoding closely related FGF Receptors, FGF R1-4, are known. All four genes for FGF Rs encode Proteins with an N-terminal signal peptide, three immunoglobulin (Ig)-like domains, an acid-box region containing a run of acidic residues between the IgI and IgII domains, a transmembrane domain and the split tyrosine-kinase domain. Multiple forms of FGF R1-3 are generated by alternative splicing of the mRNAs. A frequent splicing event involving FGF R1 and 2 results in Receptors containing all three Ig domains, referred to as the  $\alpha$  isoform, or only IgII and IgIII, referred to as the  $\beta$  isoform. Only the  $\alpha$  isoform has been identified for FGF R3 and FGF R4. Additional splicing events for FGF R1-3, involving the C-terminal half of the IgIII domain encoded by two mutually exclusive alternative exons, generate FGF Receptors with alternative IgIII domains (IIIb and IIIc). The complex patterns of expression of these Receptors as well as the specificity of their interactions with the various FGF Ligand family members are under investigation.

