

A Reliable Research Partner in Life Science and Medicine

Elab Fluor® Violet 450 Anti-Human CD18 Antibody[60.3]

Catalog Number: GFH1412Q

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Host Mouse

Isotype Mouse IgG2a, κ

Clone No. 60.3

Isotype Control Elab Fluor® Violet 450 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product GFH09802Q]

Conjugation Elab Fluor[®] Violet 450

Conjugation Information Elab Fluor[®] Violet 450 is designed to be excited by the violet laser (405 nm) and

detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of normal human peripheral blood cells with Elab

Fluor[®] Violet 450 Anti-Human CD18 Antibody[60.3] (filled gray histogram) or Elab Fluor[®] Violet 450 Mouse IgG2a, κ Isotype Control (empty black histogram). Cells in the lymphocytes gate were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names Integrin β2 subunit;LFA -1 β subunit;β2 integrin;ITGB2;b2 integrin;beta 2 integrin;CD18

Web: www.elabscience.cn

抗体;CD18流式抗体;人CD18;人CD18抗体;人CD18流式抗体;GFH1412

Uniprot ID P05107

For Research Use Only

Elabscience®

Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Gene ID

Background

3689

CD18, also known as integrin $\beta2$ subunit, LFA-1 β subunit, and $\beta2$ integrin, is a 90 - 95 kD type I glycoprotein. CD18 non-covalently associates with CD11a, CD11b, or CD11c. CD18 is expressed on all leukocytes. CD18 and associated α chains function in the adhesion and signaling in hematopoietic cells.