A Reliable Research Partner in Life Science and Medicine

# **Recombinant CD168 Monoclonal Antibody**

catalog number: AN301465L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

**Reactivity** Human;Rat;Mouse;African green monkey **Immunogen** Recombinant human CD168 fragment

 Host
 Rabbit

 Isotype
 IgG, κ

 Clone
 A160

Purification Protein A purified


Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.


## Applications Recommended Dilution

**WB** 1:500-1:1000

**IF** 1:50 **FCM** 1:50-1:100

#### Data







Western Blot with CD168 Monoclonal Antibody at dilution of Immunofluorescent analysis of (100% Ice-cold methanol)

1: 1000. Lane 1: MCF-7, Lane 2: LNCap, Lane 3: COS-7, fixed SW840 cells using anti-CD168 Monoclonal Antibody at Lane 4: Raw264.7, Lane 5: PC-12 dilution of 1:50.

Observed-MW:84 kDa Calculated-MW:84 kDa





Flow cytometric analysis of human CD168 expression on Jurkat cells. Cells were stained with purified anti-Human CD168, then a Alexa Fluor 488-conjugated second step antibody. The histogram were derived from events with the forward and side light-scatter characteristics of intact cells.

Flow cytometric analysis of human CD168 expression on K562 cells. Cells were stained with purified anti-Human CD168, then a Alexa Fluor 488-conjugated second step antibody. The histogram were derived from events with the forward and side light-scatter characteristics of intact cells.

Rev. V1.0

#### **Preparation & Storage**

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

#### **Background**

### For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: w w w .elabscience.com
 Email: techsupport@elabscience.com



## **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

Receptor for hyaluronic acid (HA). Involved in cell motility. When hyaluronan binds to HMMR, the phosphorylation of a number of proteins, including PTK2/FAK1 occurs. May also be involved in cellular transformation and metastasis formation, and in regulating extracellular-regulated kinase (ERK) activity. May act as a regulator of adipogenesis.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.0