

**IL-8/CXCL8 (C-6His), Human, Recombinant**

Cat. No. : PCK205

**General Information**

<b>Synonyms</b>	Interleukin-8;IL-8;C-X-C Motif Chemokine 8;CXCL8;Emoctakin;Granulocyte Chemotactic Protein 1;GCP-1;Monocyte-Derived Neutrophil Chemotactic Factor;MDNCF;Monocyte-Derived Neutrophil-Activating Peptide;MONAP;Neutrophil-Activating Protein 1
<b>Species</b>	Human
<b>Expression host</b>	Human Cells
<b>Sequence</b>	Glu21-Ser99
<b>Accession</b>	P10145
<b>Tag</b>	C-6His
<b>Mol mass</b>	10.1 kDa
<b>Expiration date</b>	12 months

**Product feature**

<b>Purity</b>	> 95% as determined by reducing SDS-PAGE.
<b>Endotoxin (EU/μg)</b>	< 0.1
<b>Storage</b>	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
<b>Shipping</b>	Ice bag
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Background**

Interleukin-8 (IL-8) belongs to the neutrophil-specific CXC family of Chemokines. It is one of the initial Cytokines released from a variety of cell types, including T cells, endothelial cells and fibroblasts, in response to an inflammatory stimulus and acts by recruiting neutrophils, T-cells and basophils to the site of inflammation. Elevated Interleukin-8 levels are associated with the onset of a variety of disease states.